

What is Manual Lymphatic Drainage

Manual Lymphatic Drainage is designed to reroute the lymphatic system using vessels as opposed to nodes and bypassing any obstructions. It generally takes 21 days of daily treatment to achieve this endeavour. Once discharged from therapy, the pump therapy is relied on to continue moving the fluid through the channels formed through MLD.

Can a patient still receive a pump if they have CHF (Congestive Heart Failure?)

Of course, active CHF is a contraindication for a compression pump. However, some Doctors will prescribe compression therapy for a patient with a history of Congestive Heart Failure, not active CHF. Your question to the physician would be, is the patient controlled? If the patient is controlled, compression therapy is beneficial. Remember to have the patient in a sitting position while using the compression therapy pump. Teach the patient the signs and symptoms of any problems, shortness of breath, chest pain etc. Should they experience any of these symptoms they should immediately stop their therapy and contact their physician.

Can a patient with a DVT (Deep Vein Thrombosis) receive a pump?

A patient with an active DVT is not an appropriate candidate for a compression pump.

It is always a good idea to make the physician aware that an active DVT is a contraindication. Request a copy of the last Doppler (within 6 months) for your records. There has also been some question about using compression therapy with patients who have a history of DVTs and has a filter. These patients are also candidates with the same conditions as stated above.

The only difference would be the physician will generally use less compression for treatment. Teach the patient the signs and symptoms of an active DVT. If it hurts, something is wrong, stop and contact your physician.

Can a patient use a pump over a multilayer bandage system?

Yes, a patient can use a pump over a multilayer bandage system.



For more information about the Bio Compression therapy systems, please contact;

Obex Medical on 0800 656 239 or send your enquiry to sales@obex.co.nz

OBEX

BIO COMPRESSION SYSTEMS INC.

Distributed in New Zealand by:
Obex Medical Ltd • 303 Manukau Road, Epsom, Auckland 1344
0800 656 239 • www.obex.co.nz

SEQUENTIAL GRADUATED COMPRESSION PUMPS

Frequently Asked Questions

What is the importance of gradient?

Sequential pumps with calibrated gradient pressure have proved to be the best devices for reducing the lymphatic fluid from the limb in comfortable and efficient manner. The body uses various muscle groups to move the lymphatic fluid through the channels.

How do I set the pressure on the pump?

Suggested protocols have been provided with your pump, follow these suggestions unless otherwise directed by your physician or therapist. Never increase the pressure unless told so by your physician or therapist. Should you reach a plateau in your treatment and feel as though you are not achieving further reduction, decrease the pressure and monitor. It is possible that you may be using too much pressure.

We **connect** New Zealand
with **world leaders** in medical
technology

OBEX

How long do I use the pump?

The suggested protocols provided with your pump are designed to help you achieve the maximum benefit of your therapy. Again, you should follow these instructions unless otherwise directed by your physician or therapist. In some severe cases the physician and/or therapist may deviate from the provided protocols based on your condition.

What is the application of the pump for treating ulcers and open wounds?

The pump can be used on patients with venous ulcers and/or open wounds as long as there are no signs of an untreated infection/cellulitis. If you have cellulitis you can use the pump once antibiotics have been administered for at least 72 hours.

Signs and symptoms for infection are: heat (hot extremity), rash, red streaks, pain and/or fever etc. If one or more of these symptoms occur while under compression therapy treatment discontinue your treatment and contact your physician. Wounds must be covered with the appropriate dressings and it is recommended that daily dressing changes be ordered. You should clean and change your dressing immediately following your treatment with the pump. The wound will drain significantly during pump therapy and this is a good sign.

Can the pump be used on pregnant women?

Some pregnant women will develop lymphoedema lasting the entire term of their pregnancy, but it disappears when the baby is born. However, others can continue to have a problem with oedema even after delivery. The patient should wear compression pantyhose and elevate their limbs as often as possible. Using a pump during pregnancy presents no adverse effects. The pressures for the pump are set according to the protocols used for primary lymphoedema.

How are compression garments used in treating Lymphoedema?

Compression garments are specially designed to maintain and support the limb, not to reduce its size. Garments must be applied in the morning to prevent gravity from pulling the fluid down into the limb. If this happens the garment will trap the fluid and the garments will not fit comfortably.

The patient should use rubber gloves to help in the application of the garment, as these will provide resistance. Garments should be worn daily and removed at night. Compression garments are available in standard and custom styles. There are several companies to choose from with the final choice usually coming down to cosmetics and cost. Garments usually last about 6 months, at which time it is necessary to be re-fitted. The patient should be re-measured each time a new garment is ordered to account for any changes in the size of the limb (larger or smaller). The garments are machine washable but seem to last longer when hand washed.

The patient will know it is time to get re-fitted when they notice that their limb tends to swell slightly at the end of the day or they notice any signs of stretching in the garment fabric.



How does bandaging or wrapping control Lymphoedema?

This treatment utilizes a four-layer wrap to work in conjunction with the pump and compression garments in controlling oedema.

All the digits (fingers and toes) are wrapped individually. The hand or foot is then wrapped, followed by the forearm/calf, elbow/knee, upper arm and thigh. This therapy can be taught to some patients, however older patients or those with limited dexterity i.e. arthritis, etc. may have difficulty applying these garments. The bandages are reusable.

How important is hygiene for Lymphoedema patients?

Meticulous skin care especially with the oedematous limb is essential. Because of the increased fluid levels under the skin, the skin cannot resist rips and tears in the same manner as non-oedematous skin. Any breakdown in the integrity of the skin results in susceptibility to bacteria, infection and cellulitis, the most serious of complications facing the patient.

Treat all cuts, burns and bruises, hangnails, ingrown toenails, ingrown hairs, razor rashes, blisters, scrapes, mosquito bites, etc., as potential sites for infection.

Should you notice any signs or symptoms of infection or cellulitis, contact your doctor immediately. Delay in treatment will enable the infection to spread to other areas throughout the body.